

Knowledge Attitudes And Practices Study

This two-volume set presents a course outline, in modular format, that deals step-by-step with the development of a Health Systems Research (HSR) proposal and field testing (Part 1) and with data analysis and report writing (Part 2).

The main purpose of this study was to assess knowledge, attitude and practices among young adults aged 18-25 years towards Leptospirosis within highly endemic postal zones in Kingston and St. Andrew, Jamaica.

Master's Thesis from the year 2017 in the subject Women Studies / Gender Studies, Hawassa University, language: English, abstract: Female Genital Mutilation (FGM) is recognized internationally as a violation of human rights of girls and women constituting an extreme form of gender discrimination with documented health consequences. The aim of this study was to assess knowledge, attitude and practice of women towards FGM practice. A community based cross-sectional study design was applied. Both quantitative and qualitative were employed. A total of 278 women at reproductive age (15-49) were sampled for the study from six randomly selected kebeles of Angacha woreda. The survey data was analyzed by SPSS soft ware version 20. Descriptive statistics such as mean, percentage and frequency were used for analyzing data. Binary Logistic Regression Model was used to analyze determinant Associated Factors regarding FGM. In addition qualitative data were analyzed thematically and the result was presented in narration. The result showed that 55.4% of women had good knowledge about FGM practice however 44.6% had poor knowledge. The majority of the women (50.4%) had negative attitude while 49.6% of them had positive attitude towards FGM practice. Tradition and fear of marriage were the major reasons for the continuation of this practice. From study participants 92.4%of responded that FGM was being practiced in their community and 77.7% of participants were undergone themselves. This study revealed that 79.5% of women were encountered physical health problems related to FGM and 59.7% of women were encountered psychological and sexual problems related to FGM. Bivariate and Multivariate analyses were carried out to identify determinant factors of practice. Age, family income, knowledge and attitude were significantly associated factors for FGM practice. [...]

Food-borne diseases are an important cause of morbidity and mortality, and a significant impediment to socioeconomic development worldwide, but the full extent and burden of unsafe food is unknown. Precise information on the burden of food-borne diseases can adequately inform policy-makers, allowing them to allocate appropriate resources for food safety control and intervention efforts. In 2007, the World Health Organization (WHO) listed food safety as an increasingly important public health issue, hence governments all over the world are intensifying their efforts to improve food safety. A study conducted by WHO to estimate the global burden of food-borne diseases revealed that the most frequent cause of food-borne illnesses were diarrheal disease agents, particularly norovirus and campylobacter spp. Food-borne diarrheal disease agents caused 230,000 deaths, particularly non-typhoidal salmonella enterica that causes diarrheal and invasive diseases. Other major causes of food-borne deaths were salmonella enterica, salmonella typhi, taenia solium, hepatitis A virus and aflatoxin. Forty percent of the food-borne disease burden was amongst children under five years of age (WHO, 2015). To prevent food-borne illness, it is necessary to understand how food becomes unsafe and what proactive measures can be taken to ensure food safety. In this context, FAO commissioned Al Markaz for Development and Marketing Consultancies (Al Markaz) to conduct this study to identify Palestinian households' knowledge, attitudes and practices (KAP) on food safety and KAP gaps as well as to develop a proper educational program for households. This report presents the main findings of the study, based on a survey of a representative sample of household consumers in the

Access Free Knowledge Attitudes And Practices Study

West Bank and Gaza Strip (WBGS).

Knowledge, Attitude and Practice (KAP) of Women towards Female Genital Mutilation GRIN Verlag

[Copyright: 6fbb5a66a1cf36fa3877f3c33821cbc8](#)